

CHAPS II: Experimental Planning



Larry Berg¹, Carl Berkowitz¹, John Ogren², Chris Hostetler³, and Rich Ferrare³



- ¹ Pacific Northwest National Laboratory
- ² NOAA Earth System Research Laboratory
- ³ NASA Langley Research Center





Outline

- General sampling strategy
- Instrument lists
- Flight Patterns
- Sampling issues
 - Number of clouds
 - Time in cloud
 - Identification of air that has been processed by clouds







General Sampling Strategy

- Optical properties: Dual air stream
 - Two inlets used during cloud penetration
 - Counter Flow Virtual Impactor (CVI)
 - Selectively samples cloud drops
 - Standard inlet
 - Samples all aerosols
 - Identical instrumentation
 - Fast response, ~1 Hz
- Aerosol composition: Standard inlet
 - Slower response instruments
 - Detailed size and composition
- NASA King Air flying above the G-1





Sampling Issues

- Statistical approach
 - Sample clouds along a horizontal leg
 - No need to sample the same cloud repeatedly
- Two related issues
 - How many clouds?
 - How long in a single cloud?
- How to identify air that has been processed








- LES of cloud field
 - 128 x 128 x 128 grid
 - $-\Delta X = \Delta Y = 50 \text{ m}$
 - $-\Delta Z = 40 \text{ m}$
- Fair weather GEWEX GCSS study (Brown et al. 2001)
 - Based on conditions from ARM Central Facility





LES Cloud Field



Z = 1.3 km

Cloud fraction = 10%






Sample Histograms

- Track cloud chord lengths along each leg
 - Used two time periods ~ 70 km long leg

Z (km)	# Chords	# Chords > 0.3 km
1.3	9	5
1.5	16	5
1.7	8	4
1.8	13	4








How Long in a Cloud

- Cloud Chord Length (CCL) can be determined from instruments at ARM Central Facility
 - Cloud radar, lidar, radar wind profiler
 - Samples cloud field like an aircraft
- Mean CCL ~1 km
- Approximately 60% of observed chord lengths greater than 0.5 km



From Berg and Kassianov (2005)



Identification of Cloud Processed Air




- Thermodynamic tracer of opportunity
- Variables that are conserved for dry and moist adiabatic processes
 - As air parcels rise, these values do not change
 - Equivalent potential temperature (θ_e)
 - $\theta_e = \theta \exp[L_v r_s / (C_p T)]$
 - Total water mixing ratio (r_{Total})
 - Water vapor + cloud water
- Similar analysis has been done looking at mixing in clouds






LES Sample Profiles







SP Sample LES Mixing Diagram




Z = 1.3 km






SP Sample LES Mixing Diagram







Sample LES Clouds







Conclusions


- Instrument list
- Flight pattern
- Issues
 - Number of clouds
 - Length of time in clouds
 - Identification of air that has been processed by clouds



Flight Pattern: A Statistical Study



- G-1
 - Stacks of three legs
 - Below, in and above clouds
- NASA King Air
 - Single altitude above G-1
- Preliminary okay from FAA






Instrument List: Dual Air Stream



- Aerosol scattering; Nephelometer
- Aerosol absorption; PSAP, LAPA
- CCN counters
 - Two per inlet

- Time-resolved aerosol collector
 - Single particle collection
- Water Vapor
- Aerosol size; Fast integrated mobility spectrometer
- Particle number










- Hygroscopicity; H-FIMES
- Size
 - SMPS; 0.1 3 μm
 - PCASP; 0.17-3 μm
 - CAPS; 0.5-50 μm

- EC Mass
 - DMT Particle Soot Photometer
- Bulk composition; AMS
- Particle ionic composition; PILS









- CO
- SO₂O₃
- - NOAA B2B System
- H₂O₂